

Propagation of hydrodynamic interactions between particles in a compressible fluid

Rei Tatsumi* and Ryoichi Yamamoto†

Department of Chemical Engineering, Kyoto University, Kyoto 615-8510, Japan

(Dated: November 20, 2012)

Hydrodynamic interactions are transmitted by viscous diffusion and sound propagation: the temporal evolution of hydrodynamic interactions by both mechanisms is studied by direct numerical simulation in this paper. The hydrodynamic interactions for a system of two particles in a fluid are estimated by the velocity correlation of the particles. In an incompressible fluid, hydrodynamic interactions propagate instantaneously at the infinite speed of sound, followed by the temporal evolution of viscous diffusion. On the other hand, in a compressible fluid, sound propagates at a finite speed, which affects the temporal evolution of the hydrodynamic interactions by the order of magnitude relation between the time scales of viscous diffusion and sound propagation. The hydrodynamic interactions are characterized by introducing the ratio of these time scales as an interactive compressibility factor.

I. INTRODUCTION

In particle dispersions, the motion of each particle in a fluid solvent affects the motion of the other particles. Such dynamical interactions are called hydrodynamic interactions; these interactions produce complex dynamical behavior for dispersions, which can be observed in particle aggregation and sedimentation phenomena.

Here, hydrodynamic interactions correspond to momentum exchange among particles through the ambient fluid. Hydrodynamic interactions are transmitted by two mechanisms: viscous diffusion and sound propagation. These two mechanisms occur at different time scales. The time scale for viscous diffusion over a distance equal to the particle size is $\tau_\nu = a^2/\nu$, while the time scale for sound propagation is $\tau_c = a/c$, where a is the particle radius, ν is the kinematic viscosity, and c is the speed of sound in the fluid. To study dynamical effects at the particle size scale, the relative significance of the sound propagation mechanism in hydrodynamic interactions can be assessed from the ratio of the two time scales described above:

$$\varepsilon = \frac{\tau_c}{\tau_\nu} = \frac{\nu}{ac}. \quad (1)$$

This dimensionless quantity is called the compressibility factor. Because sound propagation is much faster than viscous diffusion, the compressibility factor is generally quite small; for example, in a dispersion of water and particles of radius $a = 100$ nm, the compressibility factor is estimated to be $\varepsilon \approx 7 \times 10^{-3}$. In this case, the assumption of incompressibility is fully justified. Thus, fluids are assumed to be incompressible in many theoretical studies of hydrodynamic interactions and viscous diffusion is only considered in temporal evolution mechanisms for hydrodynamic interactions [1, 2]. On the other hand, the speed of sound in a liquid is around 10^3 m/s

irrespective of the liquid considered, producing a large compressibility factor for a dispersion of a highly viscous fluid solvent; for example, $\varepsilon \approx 0.7$ for olive oil and $\varepsilon \approx 10$ for corn syrup.

In recent years, the temporal evolution of hydrodynamic interactions between two particles has been directly observed [3–6]. In these experimental studies, particles were trapped by optical tweezers and the correlations between the positional fluctuations of particles were measured. The authors reported the temporal evolution of hydrodynamic interactions in the viscous diffusion regime, which coincided with analytical predictions that assumed the fluid was incompressible. Evidence for hydrodynamic interactions by sound propagation was also observed [3, 7, 8], but could not be captured due to the extremely short time scale of sound propagation.

In the present study, we investigate the temporal evolution of hydrodynamic interactions by modeling sound propagation using direct numerical simulation. Within this approach, the hydrodynamic interactions are directly computed directly by simultaneously solving for the motions of the fluid and the particles with appropriate boundary conditions. We use the smoothed profile method (SPM) [9, 10], which can be applied to a compressible fluid as well as an incompressible fluid [11]. Sound propagation can only be captured by accounting for fluid compressibility.

We consider a system of two particles in a fluid and investigate the correlated motion of the particles. In particular, we estimate the velocity relaxation of the particle in response to the exertion of an impulsive force on the other particle. This velocity cross-relaxation function is equivalent to the velocity cross-correlation function from the fluctuation-dissipation theorem. The numerical results for the velocity cross-relaxation function may be interpreted in terms of the temporal evolution of the flow field around the particle. We first consider an incompressible fluid to identify the separate characteristics of sound propagation and viscous diffusion in the hydrodynamic interactions. We then consider a compressible fluid to consider the effect of the order of magnitude relation of the sound propagation and viscous diffusion time

* tatsumi@cheme.kyoto-u.ac.jp

† ryoichi@cheme.kyoto-u.ac.jp

scales on the temporal evolution of the hydrodynamic interactions. In addition, we examine the validity of analytical solutions within the Oseen approximation, which are often compared with experimental results.

II. MODEL

A. Basic equations

We obtain solutions to describe the dynamics of particles in a Newtonian fluid. The motion of the particles is governed by Newton's and Euler's equations of motion, which can be written for the i -th particle as

$$M_i \frac{d}{dt} \mathbf{V}_i = \mathbf{F}_i^H + \mathbf{F}_i^C + \mathbf{F}_i^E, \quad \frac{d}{dt} \mathbf{R}_i = \mathbf{V}_i, \quad (2)$$

$$\mathbf{I}_i \cdot \frac{d}{dt} \boldsymbol{\Omega}_i = \mathbf{N}_i^H + \mathbf{N}_i^E, \quad (3)$$

where $\mathbf{R}_i$, $\mathbf{V}_i$, and $\boldsymbol{\Omega}_i$ are the position, the translational velocity, and the rotational velocity of the i -th particle, respectively. The particle has a mass M_i and a moment of inertia $\mathbf{I}_i$. The fluid exerts a hydrodynamic force $\mathbf{F}_i^H$ and a torque $\mathbf{N}_i^H$ on the particle, while a force $\mathbf{F}_i^C$ is exerted through direct interactions among the particles. A force $\mathbf{F}_i^E$ and torque $\mathbf{N}_i^E$ are externally applied. The hydrodynamic force and torque are evaluated from the momentum conservation between the particle and the fluid.

The fluid dynamics are governed by the following hydrodynamic equations:

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0, \quad (4)$$

$$\frac{\partial \rho \mathbf{v}}{\partial t} + \nabla \cdot (\rho \mathbf{v} \mathbf{v}) = \nabla \cdot \boldsymbol{\sigma} + \mathbf{f}^R, \quad (5)$$

where $\rho(\mathbf{r}, t)$ and $\mathbf{v}(\mathbf{r}, t)$ are the mass density and velocity fields, respectively, of the fluid. The stress tensor is given by

$$\boldsymbol{\sigma} = -p\mathbf{I} + \eta[\nabla \mathbf{v} + (\nabla \mathbf{v})^T] + \left(\eta_v - \frac{2}{3}\eta\right)(\nabla \cdot \mathbf{v})\mathbf{I}, \quad (6)$$

where $p(\mathbf{r}, t)$ is the pressure, η is the shear viscosity, and η_v is the bulk viscosity. A body force $\mathbf{f}^R(\mathbf{r}, t)$ is added to satisfy particle rigidity. We also assume a barotropic fluid described by $p = p(\rho)$ with a constant speed of sound c such that

$$\frac{dp}{d\rho} = c^2. \quad (7)$$

Equations (4)-(7) are closed for the variables ρ , $\mathbf{v}$, and p , without consideration of energy conservation.

We use SPM to perform direct numerical simulations. In this method, the boundaries between particle and fluid are modeled by a continuous interface. For this purpose,




FIG. 1. Geometry of the present model system. Two spherical particles of equal radius a are situated in a fluid with a center-to-center distance $R = |\mathbf{R}|$. This system is axisymmetric about the $\mathbf{R} = \mathbf{R}_2 - \mathbf{R}_1$ direction, such that two directions are of interest: parallel and perpendicular to the center-to-center axis of the particles. In this figure, the y and z directions are degenerate perpendicular directions.

a smoothed profile function $\phi(\mathbf{r}, t) \in [0, 1]$ is introduced to distinguish between the particle and fluid domains: i.e., $\phi = 1$ in the particle domain and $\phi = 0$ in the fluid domain. These two domains are smoothly connected through a thin interfacial region of thickness ξ . The body force for the particle rigidity $\mathbf{f}^R$ is expressed as $\rho\phi\mathbf{f}_p$. The detailed mathematical expressions of ϕ and $\phi\mathbf{f}_p$ have been previously given [9].

B. Linear formulation

The hydrodynamic equations can be linearized for slow flows in a dispersion. Correspondingly, the equations of motion of the particles are also linear. Most approximate theories are based on this formulation.

Here, for a N_p -particle system, we define the following column vectors to describe the equations concisely:

$$\mathbf{U} = \begin{pmatrix} \mathbf{V}_1 \\ \vdots \\ \mathbf{V}_{N_p} \\ \boldsymbol{\Omega}_1 \\ \vdots \\ \boldsymbol{\Omega}_{N_p} \end{pmatrix}, \quad \mathbf{H} = \begin{pmatrix} \mathbf{F}_1^H \\ \vdots \\ \mathbf{F}_{N_p}^H \\ \mathbf{N}_1^H \\ \vdots \\ \mathbf{N}_{N_p}^H \end{pmatrix}, \quad \mathbf{E} = \begin{pmatrix} \mathbf{F}_1^E \\ \vdots \\ \mathbf{F}_{N_p}^E \\ \mathbf{N}_1^E \\ \vdots \\ \mathbf{N}_{N_p}^E \end{pmatrix},$$

which are the $6N_p$ -dimensional vectors. Neglecting direct particle interactions, the equations of particle motion, i.e., Eqs. (2) and (3) are summarized by

$$\mathbf{M} \cdot \frac{d}{dt} \mathbf{U}(t) = \mathbf{H}(t) + \mathbf{E}(t), \quad (8)$$

where the matrix $\mathbf{M}$ represents the direct sum of the mass tensor and the moment of inertia tensor. For spherical particles with equal radii and densities, the matrix $\mathbf{M}$ reduces to

$$\mathbf{M} = \begin{pmatrix} M\mathbf{I}_{3N_p} & \mathbf{0} \\ \mathbf{0} & \frac{2}{5}Ma^2\mathbf{I}_{3N_p} \end{pmatrix}, \quad (9)$$

where M and a are the mass and the radius, respectively, of each particle. With Fourier transforming Eq. (8), the corresponding equation for the Fourier components with the time factor $e^{-i\omega t}$ is obtained by

$$-i\omega \mathbf{M} \cdot \hat{\mathbf{U}}(\omega) = \hat{\mathbf{H}}(\omega) + \hat{\mathbf{E}}(\omega). \quad (10)$$

The following relation holds true for the linearized hydrodynamic equations [2]:

$$\hat{\mathbf{U}}(\omega) = -\hat{\boldsymbol{\mu}}(\omega) \cdot \hat{\mathbf{H}}(\omega), \quad (11)$$

where $\hat{\boldsymbol{\mu}}$ is the $6N_p \times 6N_p$ mobility matrix. The mobility matrix depends on the configuration of all the particles $\{\mathbf{R}_i\}$, so we further assume, for simplicity, that the particles do not move significantly over the time scale considered, namely, the configuration of particles $\{\mathbf{R}_i\}$ is independent of time t . Substitution of Eq. (11) into Eq. (10) yields an explicit expression for the translational and rotational particle velocities:

$$\hat{\mathbf{U}}(\omega) = [\mathbf{I} - i\omega \hat{\boldsymbol{\mu}}(\omega) \cdot \mathbf{M}]^{-1} \cdot \hat{\boldsymbol{\mu}}(\omega) \cdot \hat{\mathbf{E}}(\omega). \quad (12)$$

Therefore, for an externally applied force and torque, the dynamics of particles in a fluid can be described by the mobility matrix.

The components of the mobility matrix are given by

$$\hat{\boldsymbol{\mu}}(\omega) = \begin{pmatrix} \hat{\boldsymbol{\mu}}^{\text{tt}}(\omega) & \hat{\boldsymbol{\mu}}^{\text{tr}}(\omega) \\ \hat{\boldsymbol{\mu}}^{\text{rt}}(\omega) & \hat{\boldsymbol{\mu}}^{\text{rr}}(\omega) \end{pmatrix}, \quad (13)$$

where $\hat{\boldsymbol{\mu}}^{\alpha\beta}$ is the $3N_p \times 3N_p$ matrix, which is composed of the mobility tensors $\hat{\boldsymbol{\mu}}_{ij}^{\alpha\beta}$ for two particles i and j . The mobility tensors describe the mutual coupling of the translational and rotational motion between particles and the superscript represents the mode of the motion: 't' for translation and 'r' for rotation. The mobility matrix is symmetric as a consequence of the Lorentz reciprocal theorem, such that the following relations are satisfied [12]:

$$\hat{\boldsymbol{\mu}}^{\text{tt}} = (\hat{\boldsymbol{\mu}}^{\text{tt}})^T, \quad \hat{\boldsymbol{\mu}}^{\text{rr}} = (\hat{\boldsymbol{\mu}}^{\text{rr}})^T, \quad \hat{\boldsymbol{\mu}}^{\text{tr}} = (\hat{\boldsymbol{\mu}}^{\text{rt}})^T. \quad (14)$$

Now, we consider a system of two identical spherical particles in a fluid. The configuration of the particles is described in Fig. 1. Numbers are assigned to the particles: 1 for the particle on the left and 2 for the particle on the right. We define a vector $\mathbf{R} = \mathbf{R}_2 - \mathbf{R}_1$ to describe the geometry of this system. The particle center-to-center distance is denoted by $R = |\mathbf{R}|$. The unit vector along the line of centers is described by $\hat{\mathbf{R}} = \mathbf{R}/R$. Due to axisymmetry about the $\hat{\mathbf{R}}$ axis, each mobility tensor is described by at most two scalar functions [13, 14]:

$$\hat{\boldsymbol{\mu}}_{ij}^{\text{tt}}(\mathbf{R}, \omega) = \hat{\mu}_{ij}^{\text{tt}\parallel}(R, \omega) \hat{\mathbf{R}} \hat{\mathbf{R}} + \hat{\mu}_{ij}^{\text{tt}\perp}(R, \omega) (\mathbf{I} - \hat{\mathbf{R}} \hat{\mathbf{R}}), \quad (15a)$$

$$\hat{\boldsymbol{\mu}}_{ij}^{\text{rr}}(\mathbf{R}, \omega) = \hat{\mu}_{ij}^{\text{rr}\parallel}(R, \omega) \hat{\mathbf{R}} \hat{\mathbf{R}} + \hat{\mu}_{ij}^{\text{rr}\perp}(R, \omega) (\mathbf{I} - \hat{\mathbf{R}} \hat{\mathbf{R}}), \quad (15b)$$

$$\hat{\boldsymbol{\mu}}_{ij}^{\text{tr}}(\mathbf{R}, \omega) = \hat{\mu}_{ij}^{\text{tr}\perp}(R, \omega) \hat{\mathbf{R}} \times \mathbf{I}. \quad (15c)$$

The superscripts $\parallel$ and $\perp$ denote the directions of motion parallel and perpendicular to the symmetry axis, respectively. In the direction parallel to the symmetry axis, the translational and rotational motions are decoupled as indicated by Eq. (15c). Interchanging the particle numbers 1 and 2 corresponds to inverting the direction of $\hat{\mathbf{R}}$, which causes sign inversion for the translation-rotation cross-mobility tensor $\hat{\boldsymbol{\mu}}_{ij}^{\text{tr}}$ without changing sign of the other cross-mobility tensors $\hat{\boldsymbol{\mu}}_{ij}^{\text{tt}}$ and $\hat{\boldsymbol{\mu}}_{ij}^{\text{rr}}$. The mobility tensors Eqs. (15) within the Oseen approximation are described in the Appendix.

C. Temporal evolution of the flow field

We first consider a particle dispersion with an incompressible fluid solvent. Incompressibility corresponds to an infinite speed of sound in the fluid, such that the solenoidal condition is imposed on the fluid velocity field from Eqs. (4) and (7) as

$$\nabla \cdot \mathbf{v} = 0. \quad (16)$$

The velocity field can be decomposed into two contributions as

$$\mathbf{v} = \mathbf{w} - \nabla \varphi. \quad (17)$$

The vector field $\mathbf{w}(\mathbf{r}, t)$ is the juxtaposition of the velocity fields in the fluid and particle domains, so that the fluid-particle impermeability boundary condition, or the continuity of the normal velocity on the boundary, is not satisfied. An irrotational flow field, described by the scalar potential $\varphi(\mathbf{r}, t)$, is added in order that the velocity field $\mathbf{v}$ satisfies the boundary condition. Due to the solenoidal condition on the velocity field $\mathbf{v}$, the scalar potential φ obeys the Poisson equation:

$$\nabla^2 \varphi = \nabla \cdot \mathbf{w}. \quad (18)$$

On the right-hand side, $\nabla \cdot \mathbf{w}$ is zero except at the fluid-particle boundary, at which singularities exist. From Eq. (18), the scalar potential is given by [15]

$$\varphi(\mathbf{r}, t) = -\frac{1}{4\pi} \int d\mathbf{r}' \frac{(\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3} \cdot \mathbf{w}(\mathbf{r}', t). \quad (19)$$

Now we consider an isolated single particle, due to axisymmetry, the scalar potential and the corresponding velocity field can be simply described as

$$\varphi(\mathbf{r}, t) = \frac{Q(t) \cdot \mathbf{r}}{r^3}, \quad (20)$$

$$-\nabla \varphi(\mathbf{r}, t) = \frac{Q(t) \cdot (3\hat{\mathbf{r}}\hat{\mathbf{r}} - \mathbf{I})}{r^3}, \quad (21)$$

where $\hat{\mathbf{r}} = \mathbf{r}/r$ is a unit vector in the radial direction. The velocity field of Eq. (21) represents a doublet flow, which corresponds to the electric field generated by an electric

dipole. The vector $\mathbf{Q}(t)$ is parallel to the particle velocity and depends on time in a reflection of the particle motion relative to the fluid. The analytical form of $\mathbf{Q}(t)$ has been derived previously [16], and the strength $Q(t) = |\mathbf{Q}(t)|$ is time-independent for a neutrally buoyant particle. The doublet flow is generated instantaneously to satisfy the solenoidal condition, thus, the doublet flow is interpreted to expand at the infinite speed of sound.

The vector field $\mathbf{w}$ represents the shear flow due to viscous diffusion. For sufficiently slow fluid flow, the hydrodynamic equations can be linearized as follows:

$$\rho_0 \frac{\partial \mathbf{v}}{\partial t} = -\nabla p + \eta \nabla^2 \mathbf{v} + \mathbf{f}^R. \quad (22)$$

Under the incompressibility condition, the pressure gradient imposes fluid-particle impermeability on the time-derivative of the velocity field, so that the pressure is related to the scalar potential as

$$p = \rho_0 \frac{\partial \varphi}{\partial t}. \quad (23)$$

Therefore, the temporal evolution of the vector field $\mathbf{w}$ is given by

$$\rho_0 \frac{\partial \mathbf{w}}{\partial t} = \eta (\nabla^2 \mathbf{w} - \nabla \nabla \cdot \mathbf{w}) + \mathbf{f}^R. \quad (24)$$

Modifying the diffusion term on the right-hand side to be solenoidal, results in the doublet flow being gradually offset by the diffusion of the vector field $\mathbf{w}$.

In short, the exertion of an impulsive force on a particle in an incompressible fluid instantaneously generates a doublet flow. The shear flow then spreads out diffusely, cancelling out the doublet flow. This picture will be validated in one of the following numerical simulation results.

For a compressible fluid, a second contribution is added to Eq. (17):

$$\mathbf{v} = \mathbf{w} - \nabla \psi - \nabla \varphi, \quad (25)$$

where the scalar potential $\psi(\mathbf{r}, t)$ corresponds to the bulk compression flow. Here, we assume that the dynamics of fluid can be described by the linearized hydrodynamic equations:

$$\frac{\partial p}{\partial t} = -\rho_0 c^2 \nabla \cdot \mathbf{v}, \quad (26)$$

$$\rho_0 \frac{\partial \mathbf{v}}{\partial t} = \eta \nabla^2 \mathbf{v} + \left(\frac{1}{3} \eta + \eta_v \right) \nabla \nabla \cdot \mathbf{v} - \nabla p + \mathbf{f}^R. \quad (27)$$

The time evolution equation of the total scalar potential $\varphi' = \varphi + \psi$ is derived from the equations above as

$$\begin{aligned} \nabla^2 \varphi' - \frac{1}{c^2} \frac{\partial^2 \varphi'}{\partial t^2} + \frac{\eta_l}{\rho_0 c^2} \nabla^2 \frac{\partial \varphi'}{\partial t} \\ = \left(1 + \frac{\eta_l}{\rho_0 c^2} \nabla^2 \frac{\partial}{\partial t} \right) (\nabla \cdot \mathbf{w}), \end{aligned} \quad (28)$$

where $\eta_l = (4/3)\eta + \eta_v$ denotes the longitudinal viscosity. This is a damped wave equation, where the source is given on the right-hand side of the equation. Therefore, a potential flow propagates at a finite speed of sound in a compressible fluid. In the long-time limit, Eq. (28) reduces to the Poisson equation for an incompressible fluid, as given by Eq. (18), and the potential flow reduces to an instantaneous doublet flow.

III. NUMERICAL RESULTS

Numerical simulations are performed for a three-dimensional box with periodic boundary conditions. The space is divided into meshes of length Δ , which is a unit length. The units of the other physical quantities are defined by combining $\eta = 1$ and $\rho_0 = 1$ with Δ , where ρ_0 is the fluid mass density at equilibrium. The system size is $L_x \times L_y \times L_z = 256 \times 256 \times 256$. The other parameters are set to $a = 4$, $\xi = 2$, $\rho_p = 1$, $\eta_v = 0$, and $h = 0.05$, where ρ_p is the particle mass density and h is the time increment of a single simulation step.

We consider a system in which two identical spherical particles in a fluid, whose geometry is described in Fig 1. Both particles have the same density and radius, namely the same mass and moment of inertia. We investigate the time-dependence of the velocity for particle 1 following the exertion of an impulsive force at the center of particle 2, with changing the center-to-center distance between particles given by $R^* = R/a$. This cross-relaxation function is a manifestation of the temporal evolution of the hydrodynamic interactions between particles 1 and 2. The impulsive force is assumed to be sufficiently small such that the Reynolds and Mach numbers of the flow are sufficiently low. We set the impulsive force to produce an initial particle Reynolds number of $\text{Re}_p = 10^{-3}$. In addition, because the particle displacement before stopping, scaled by the particle radius a , is comparable to $(\rho_p/\rho_0)\text{Re}_p$, the displacement of particles is negligible in the present simulations. Therefore, because particle collisions do not occur, the direct interactions between particles, including overlap repulsion forces, are not considered.

The cross-relaxation tensor is introduced as the normalized change in velocity of particle 1:

$$\mathbf{V}_1(\mathbf{R}_1, t; \mathbf{R}_2) = \frac{\mathbf{P}_2}{M} \cdot \gamma_{12}(\mathbf{R}, t), \quad (29)$$

$$\gamma_{12}(\mathbf{R}, t) = \gamma_{12}^{\parallel}(R, t) \hat{\mathbf{R}} \hat{\mathbf{R}} + \gamma_{12}^{\perp}(R, t) (\mathbf{I} - \hat{\mathbf{R}} \hat{\mathbf{R}}), \quad (30)$$

where $\mathbf{P}_2$ is the impulsive force exerted on particle 2 at $t = 0$. Due to the previously mentioned axisymmetry, the cross-relaxation tensor is characterized by only two directions of motion, which are parallel and perpendicular to the center-to-center vector $\mathbf{R}$. Then, the motions in both directions are decoupled. From the fluctuation-dissipation theorem, the cross-relaxation tensor is equiv-

alent to the velocity cross-correlation function of the fluctuating system:

$$\gamma_{12}(\mathbf{R}, t) = \frac{M}{k_B T} \langle \mathbf{V}_1(\mathbf{0}, 0) \mathbf{V}_2(\mathbf{R}, t) \rangle, \quad (31)$$

where k_B is the Boltzmann constant and T is the thermodynamic temperature. We examine both the parallel correlation $\gamma_{12}^{\parallel}$ and the perpendicular correlation $\gamma_{12}^{\perp}$ by adjusting the direction of the impulsive force $\mathbf{P}_2$. As predicted by the formulation in the preceding section, for motion perpendicular to the axis, the translation-rotation coupling in the particle motion is also observed. However, because the influence of sound propagation on the rotational motion is expected to be small, we will focus entirely on the translation-translation coupling. We also compare the simulated results with approximate solutions, where the analytical solution for an isolated single particle for the self-mobilities and the Oseen approximation for the cross-mobilities are applied (see the Appendix).

A. Incompressible fluid

First, we investigate the temporal evolution of hydrodynamic interactions in an incompressible fluid. The dynamics of the incompressible fluid is governed by the hydrodynamic equations composed of Eqs. (5), (6), and (16). The center-to-center distance between particles takes values of $R^* = 3, 4$, and 7 . The simulation results for the velocity cross-relaxation functions are shown in Fig. 2. While the parallel correlation $\gamma_{12}^{\parallel}(t)$ is positive at all times, The perpendicular correlation $\gamma_{12}^{\perp}(t)$ is initially negative and subsequently becomes positive. Comparable results have been reported in experimental studies [3–5], where the difference between the parallel and perpendicular correlations has been attributed to the flow field around the particle [4, 7]. We discuss the time-dependence of the velocity cross-relaxation functions in more detail below.

The temporal evolution of the flow field has been explained in the previous section. Here, the velocity fields generated by the impulsive force exerted on a single particle are shown in Fig. 3. These simulation results are obtained for a single particle system with size $L_x \times L_y \times L_z = 128 \times 128 \times 128$. At early times $t/\tau_v = 2.5 \times 10^{-2}$, the doublet flow is dominant. The doublet flow is characterized by loop streamlines flared in the direction perpendicular to the particle motion, where backflow is observed as described by Eq. (20). As discussed in the previous section, the doublet flow appears instantaneously, which is interpreted as the propagation of an infinite-speed sound wave. On the other hand, the shear flow due to viscous diffusion, whose strength is given by the vorticity $\nabla \times \mathbf{v}$, is only observed in the vicinity of the particle. Over a time $t/\tau_v = 1.2$, the shear flow has diffused over a large range following Eq. (24); therefore, there is a spreading




FIG. 2. Velocity cross-relaxation function in an incompressible fluid in (a) parallel and (b) perpendicular directions relative to the symmetry axis of the particles. The distance between the two particles are $R^* = 3, 4$, and 7 . Simulation results are described by bold solid lines (positive values) and dashed-dotted lines (negative values). Solutions within the Oseen approximation are represented by thin solid lines (positive values) and dashed-dotted lines (negative values). The broken line represents the long-time tail with the algebraic power law decay $Bt^{-3/2}$, as given by Eq. (32).

fluid region flowing in the same direction as the particle motion, while the loop streamlines get away from the particle.

The velocity cross-relaxation function is related to the temporal evolution of the flow field around particle 2. The doublet flow propagated by sound waves produces an instantaneous velocity correlation between the particles. The doublet flow direction shown in Fig. 3 produces the positive parallel correlation and the negative perpendicular correlation. Little change in the cross-relaxation functions occurs in the early stages, reflecting the time-independence of the strength of the doublet flow due to




FIG. 3. Temporal evolution of the velocity field around an isolated single particle due to an impulsive force exerted on the particle in an incompressible fluid. The cross-sections parallel to the impulsive force direction including the particle center are shown. The impulsive force is exerted at time $t = 0$ and the simulation results are given at (a) $t/\tau_\nu = 2.5 \times 10^{-2}$ and (b) $t/\tau_\nu = 1.2$. The direction of the impulsive force is to the right in the pictures and the particle is represented by a black circle. The velocity fields are plotted only for $|\mathbf{v}|(a/\nu\text{Re}_p) > 10^{-4}$. The vorticity of the velocity field $\nabla \times \mathbf{v}$ is described by a color scale (gray scale), going from negative (darker) to positive (lighter) vorticity. The vorticity is normalized by τ_ν/Re_p .




FIG. 4. Temporal evolution of the velocity field around the particles. The cross-sections including the center-to-center axis between particles are shown. The center-to-center distance between two particles is $R^* = 7$. The impulsive force is exerted at time $t = 0$ on the particle on the right and the simulation results are presented at (a,b) $t/\tau_\nu = 6.25 \times 10^{-1}$ and (c,d) $t/\tau_\nu = 12.5$. The direction of the impulsive force is (a,c) to the right and (b,d) upwards. The particles are represented by black circles. The velocity fields are shown only for $|\mathbf{v}|(a/\nu\text{Re}_p) > 10^{-4}$. The vorticity of the velocity field $\nabla \times \mathbf{v}$ is described by a color scale (gray scale), going from negative (darker) to positive (lighter) vorticity. The vorticity is normalized by τ_ν/Re_p .




FIG. 5. Velocity cross-relaxation function in a compressible fluid in (a, b) parallel and (c, d) perpendicular directions to the symmetry axis. The center-to-center distance between two particles is $R^* = 3$. The compressibility factor takes values of $\epsilon = 0.1, 0.6$, and 1.5 . Simulation results are described by bold solid lines (positive values) and dashed-dotted lines (negative values). Solutions within the Oseen approximation are represented by thin solid lines (positive values) and dashed-dotted lines (negative values).

the neutrally buoyant particle [16].

The reduction in the parallel correlation and the sign inversion of the perpendicular correlation occur at about the same time, which depends on the inter-particle distance R^* . Because shear flow by viscous diffusion can cause such changes in the cross-relaxation functions, we estimate the time scale on which the shear flow generated by the particle 2 arrives at particle 1 by the viscous diffusion time scale over the length L : $\tau_\nu^* = L^2/\nu$. In this case, the characteristic length is the distance between the particle surfaces, $L = R - 2a$; thus, for $R^* = 3, 4$, and 7 , the viscous diffusion time scales are $\tau_\nu^*/\tau_\nu = (R^* - 2)^2 = 1, 4$, and 25 , respectively. The time for the parallel correlation to begin decreasing and for the perpendicular correlation to invert the sign roughly corresponds to τ_ν^* in each case.

In an incompressible fluid, hydrodynamic interactions are instantly propagated at the infinite speed of sound and subsequently transmitted by viscous diffusion in a time scale τ_ν^* . The temporal evolution of the flow fields around two particles separated by $R^* = 7$ is shown in Fig. 4. At early times in the flow, $t/\tau_\nu = 6.25 \times 10^{-1}$ so that the shear flow does not arrive at particle 1, whose dynamics are governed by the doublet flow. Later, at $t/\tau_\nu = 12.5$, the shear flow diffuses over time to arrive at particle 1 and the motion of particle 1 is governed by the shear flow. Here, the stresslet flow field is observed in the vicinity of particle 1. This flow is generated to maintain particle rigidity against deformation in a shear flow. In the perpendicular correlation, the rotational motion of particle 1 is also observed.

In the final stages of the flow, the momentum of parti-




FIG. 6. Velocity cross-relaxation function in a compressible fluid in (a, b) parallel and (c, d) perpendicular directions to the symmetry axis. The center-to-center distance between two particles is $R^* = 7$. The compressibility factor takes values of $\varepsilon = 0.1, 0.6$, and 1.5 . Simulation results are described by bold solid lines (positive values) and dashed-dotted lines (negative values). Solutions within the Oseen approximation are represented by thin solid lines (positive values) and dashed-dotted lines (negative values).

cle 2 has completely diffused away, so that the particles and the fluid move collectively. These dynamics are manifested in a long-time tail with a power law decay $t^{-3/2}$ of the cross-relaxation functions shown in Fig. 2, which are given by [17]

$$\gamma_{12}^{\parallel,\perp}(t) = \frac{1}{9\sqrt{\pi}} \frac{\rho_p}{\rho_0} \left(\frac{\tau_\nu}{t} \right)^{3/2} \quad \text{as } t \rightarrow \infty. \quad (32)$$

Discrepancies between the simulation results and the Oseen approximation are observed, especially at $t/\tau_\nu \lesssim 3$, as shown in Fig. 2. The approximate cross-relaxation functions clearly change with time in the early stages and the time at which the effect of the shear flow becomes apparent is later than in the simulation results. Particles are assumed to be points in the Oseen approximation; therefore, this approximation implies that the particle separation is much larger than the particle radius as

$R^* = R/a \gg 1$ and that the time scale of observation is much longer than that of viscous diffusion over the length of the particle radius as $t/\tau_\nu = t\nu/a^2 \gg 1$. Neglecting the particle size results in a larger characteristic length $L = R$ and a longer diffusion time scale $\tau_\nu^*/\tau_\nu = R^{*2}$. Consequently, the Oseen approximation can only be accurately applied at long times and large particle separations. The simulation results shown in Fig. 2 confirm that the validity of the Oseen approximation increases with increasing inter-particle distance R^* . In previous experimental studies, measurements were conducted over the range for which the Oseen approximation is valid, i.e., $R^* \gtrsim 5$ and measuring frequencies corresponding to $t/\tau_\nu \gtrsim 30$: consequently, the experimental results showed good agreement with the Oseen approximation[3, 4].




FIG. 7. Velocity field around particles in a compressible fluid. Simulation results at $t/\tau_v = 6.25 \times 10^{-1}$ for compressibility factors with values (a,b) $\varepsilon = 0.1$ and (c,d) $\varepsilon = 1.5$. The velocity fields are shown only for $|\mathbf{v}|(a/\nu\text{Re}_p) > 10^{-4}$. The divergence of the velocity field $\nabla \cdot \mathbf{v}$ is described by a color scale (gray scale), going from negative (darker) to positive (lighter) divergence. The value of divergence is normalized by τ_v/Re_p .

B. Compressible fluid

Next, we consider a compressible fluid in which sound propagates at a finite speed. The velocity cross-relaxation functions for inter-particle distances $R^* = 3$ and 7 are shown in Figs. 5 and 6, respectively. Due to the periodic boundary conditions imposed on the simulation box, a sound pulse generated by the particle motion can affect the simulation results after the sound pulse arrives at the end of the simulation box. The oscillatory structure of the cross-relaxation functions, especially at $\varepsilon = 0.1$, is one of the striking artifacts of the periodic boundary conditions.

The velocity cross-relaxation functions are zero in the early stages and then suddenly change to non-zero values. Because this time lag increases with the inter-particle separation, this is likely to be due to the sound propagation from particle 2 to particle 1. As with the diffusion of shear flow, we estimate the time scale for sound propagation over a characteristic length $L = R - 2a$ as $\tau_c^* = L/c$. For $R^* = 3$ and 7, the sound propagation time scales are $\tau_c^*/\tau_v = (R^* - 2)\varepsilon = \varepsilon$ and 5ε , respectively. The

results shown in Figs. 5 and 6 confirm the correspondence between the time scale of sound propagation and the peak of the cross-relaxation functions. The peak in the cross-relaxation function broadens as sound propagation is dissipated by the longitudinal viscosity according to Eq. (28).

When the fluid compressibility is as small as $\varepsilon = 0.1$, the velocity cross-relaxation functions superimpose onto those for the incompressible fluid after a time τ_c^* . This behavior only describes hydrodynamic interactions by sound propagation before viscous diffusion effects come into play. On the other hand, for a large compressibility $\varepsilon = 1.5$, the behavior of the cross-relaxation functions are considerably different, except for the hydrodynamic long-time tail: in particular, there is no negative perpendicular correlation. For a fluid with medium compressibility such as $\varepsilon = 0.6$, the negative perpendicular correlation is observed only for inter-particle separations as large as $R^* = 7$. As sound propagation and viscous diffusion generate negative and positive perpendicular correlations, respectively, the balance of the two time scales τ_v^* and τ_c^* is expected to characterize the behavior of the




FIG. 8. Velocity field around particles in a compressible fluid. Simulation results at $t/\tau_\nu = 3.13$ for compressibility factor with value $\varepsilon = 1.5$. The velocity fields are shown only for $|\mathbf{v}|(a/\nu\text{Re}_p) > 10^{-4}$. The (a,b) divergence and (c,d) vorticity of the velocity field are described by a color scale (gray scale), going from negative (darker) to positive (lighter) divergence. The value of divergence is normalized by τ_ν/Re_p .

cross-relaxation function. Therefore, we define the interactive compressibility factor using a characteristic length $L = R - 2a$ as

$$\varepsilon^* = \frac{\tau_c^*}{\tau_\nu^*} = \frac{\varepsilon}{R^* - 2}. \quad (33)$$

As the inter-particle distance increases, the two time scales separate to reduce the interactive compressibility factor. In the present simulations, the interactive compressibility factors for $R^* = 3$ and 7 are $\varepsilon^* = \varepsilon$ and 0.2ε , respectively. When the interactive compressibility factor is small, hydrodynamic interactions propagating at the speed of sound arrive first at the other particle in about a time τ_c^* , followed by those propagating by viscous diffusion arriving in about a time τ_ν^* . The temporal evolution is qualitatively the same as in an incompressible fluid, except for the time lag due to sound propagation at a finite speed. This situation applies to the results where a negative perpendicular correlation is observed. On the other hand, if the interactive compressibility factor is sufficiently large, the order of arrival of sound propagation and viscous diffusion should be reversed. In this situation, the perpendicular correlation is positive from

the start due to viscous diffusion and the effect of sound propagation on the hydrodynamic interactions produces a local minimum at most in the cross-correlation function. In the present studies, such results are observed when at least $\varepsilon^* \geq 0.3$ is satisfied.

The velocity fields around particles separated by $R^* = 7$ at $t/\tau_\nu = 6.25 \times 10^{-1}$ are shown in Fig. 7. The expanding doublet flow with a strength given by the divergence of the velocity field $\nabla \cdot \mathbf{v}$ are observed. For a small compressibility factor $\varepsilon = 0.1$, the doublet flow has just arrived at particle 1. This situation corresponds to a peak in the cross-relaxation functions shown in Fig. 6(a,c). On the other hand, for a large compressibility factor $\varepsilon = 1.5$, the doublet flow has not yet arrived at particle 1 and the cross-relaxation functions are correspondingly zero. At a later time $t/\tau_\nu = 3.13$, as shown in Fig. 8, the shear flow (vorticity) arrives at particle 1, instead of the doublet flow (divergence). These dynamics are manifested in the positive cross-relaxation functions of the perpendicular correlation shown in Fig. 6(b,d).

In a compressible fluid, because there is a finite sound propagation time scale as well as a viscous diffusion time

scale, the time condition for the validity of the Oseen approximation is given by $t \gg \max(\tau_\nu, \tau_c)$. As with the diffusion of shear flow, sound propagation from particle 2 to particle 1 is delayed within the Oseen approximation as shown in Figs. 5 and 6. This is because the sound propagation time scale is increased within this approximation so that $\tau_c^*/\tau_\nu = R^*\varepsilon$. Thus, the discrepancies with the simulation results can be fairly well-resolved by increasing the particle separation.

IV. CONCLUSION

In the present study, we investigated the temporal evolution of hydrodynamic interactions for a system of two particles, using SPM to perform direct numerical simulations. Fluid compressibility was considered in examining temporal evolution by sound propagation. Hydrodynamic interactions were estimated by the velocity cross-relaxation functions, which are equivalent to the velocity cross-correlation functions in a fluctuating system.

In an incompressible fluid, hydrodynamic interactions were observed to propagate instantaneously at the infinite speed of sound, along with subsequent temporal evolution by viscous diffusion. Theoretical analysis showed that a doublet flow and shear flow are generated by sound propagation and viscous diffusion, respectively. Because the cross-relaxation function reflects the characteristics of each flow field, sound propagation and viscous diffusion are associated with negative and positive perpendicular correlations between the particle velocities, respectively. Therefore, the time at which the behavior of the cross-relaxation function changes is related to the time scale of viscous diffusion over the particle separation τ_ν^* , which is only the time scale characterizing the temporal evolution of hydrodynamic interactions in an incompressible fluid.

In a compressible fluid, sound propagates between particles in a finite time, which scales as τ_c^* . The effect of the order of magnitude relation between the two time scales τ_ν^* and τ_c^* on the cross-relaxation function was observed; an interactive compressibility factor ε^* was defined as the ratio of the two time scales given by Eq. (33). In our simulation results, the reversal of the order of arrival of sound propagation and viscous diffusion at the other particle was observed at $\varepsilon^* \geq 0.3$, for which there was no negative perpendicular correlation. In this case, the hydrodynamic interactions were largely governed by viscous diffusion.

The temporal evolution of hydrodynamic interactions does not qualitatively change from that for an incompressible fluid, as long as the interactive compressibility factor is small. Only when the interactive compressibility factor is sufficiently large can differences in the temporal evolution of hydrodynamic interactions be expected. As such a situation can be realized in a highly viscous fluid, experimental validation of the results presented here is desirable.

ACKNOWLEDGEMENTS

This work was supported by KAKENHI 23244087 and the JSPS Core-to-Core Program “International research network for non-equilibrium dynamics of soft matter.”

APPENDIX: MOBILITY MATRIX IN THE OSEEN APPROXIMATION

Hydrodynamic interactions among particles in slow flows are described by the mobility matrix discussed in §2.2. However, there is no known closed-form solution even for a two-particle system. In this section, we introduce an approximate analytical form for the mobility tensors.

For slow flows, the hydrodynamic equations can be linearized as given by Eqs. (26) and (27), and the corresponding equations for the Fourier components, with the time factor $e^{-i\omega t}$, are obtained as

$$-i\omega\hat{p}_\omega + \rho_0 c^2 \nabla \cdot \hat{\mathbf{v}}_\omega = 0, \quad (\text{A1})$$

$$-i\omega\hat{\mathbf{v}}_\omega = \eta \nabla^2 \hat{\mathbf{v}}_\omega + \left(\frac{1}{3}\eta + \eta_v\right) \nabla \nabla \cdot \hat{\mathbf{v}}_\omega - \nabla \hat{p}_\omega + \hat{\mathbf{f}}_\omega^R. \quad (\text{A2})$$

The fluid velocity field generated by the body force $\mathbf{f}^R$ is expressed as

$$\hat{\mathbf{v}}_\omega(\mathbf{r}) = \int d\mathbf{r}' \hat{\mathbf{G}}(\mathbf{r} - \mathbf{r}', \omega) \cdot \hat{\mathbf{f}}_\omega^R(\mathbf{r}'), \quad (\text{A3})$$

where the Green's function is given by [18, 19]

$$\hat{\mathbf{G}}(\mathbf{r}, \omega) = \frac{1}{4\pi\eta} \left(\frac{e^{-\alpha r}}{r} \mathbf{I} + \alpha^{-2} \nabla \nabla \frac{e^{i\mu r} - e^{-\alpha r}}{r} \right), \quad (\text{A4})$$

with

$$\alpha = (-i\omega\rho_0/\eta)^{1/2}, \quad \mu = \omega/\tilde{c}, \quad (\text{A5})$$

and

$$\tilde{c} = c \left[1 - \frac{i\omega}{\rho_0 c^2} \left(\frac{4}{3}\eta + \eta_v \right) \right]^{1/2}. \quad (\text{A6})$$

For an isolated single spherical particle in a fluid, with a body force constraint $\mathbf{f}^R$ to satisfy the condition of particle rigidity, the self-mobilities can be analytically derived [18, 20, 21] as follows:

$$\hat{\mu}_{11}^{\text{tt}}(\omega) = \mu_0^{\text{t}} \frac{9}{2x^2} \frac{2x^2(1-iy) - (1+x)y^2 - x^2y^2}{(1+x)(9-9iy-2y^2) + x^2(1-iy)}, \quad (\text{A7})$$

$$\hat{\mu}_{11}^{\text{rr}}(\omega) = \mu_0^{\text{r}} \frac{3(1+x)}{3+3x+x^2}, \quad (\text{A8})$$

where $x = \alpha a$, $y = \mu a$, $\mu_0^t = (6\pi\eta a)^{-1}$, and $\mu_0^r = (8\pi\eta a^3)^{-1}$. When the speed of sound is assumed to be infinite, or $\mu = 0$, the solution for an incompressible fluid is obtained. The translation-rotation coupling does not appear in the self-mobility as $\hat{\mu}_{11}^{rt} = 0$.

To calculate the cross-mobilities, we introduce the Oseen approximation in which the particles are regarded as points. In the Oseen approximation, a particle exerts a Stokeslet (point force); therefore, according to Eq. (A3), the translation-translation cross-mobility is the Green's function itself:

$$\hat{\mu}_{12}^{tt}(\mathbf{R}, \omega) = \hat{\mathbf{G}}(\mathbf{R}, \omega). \quad (\text{A9})$$

The components of the parallel and perpendicular directions are explicitly given by

$$\hat{\mu}_{12}^{tt\parallel}(R, \omega) = \mu_0^t \frac{3a}{2\alpha^2 R^3} [(2 - 2i\mu R - \mu^2 R^2)e^{i\mu R} - (2 + 2\alpha R)e^{-\alpha R}], \quad (\text{A10a})$$

$$\hat{\mu}_{12}^{tt\perp}(R, \omega) = \mu_0^t \frac{3a}{2\alpha^2 R^3} [(-1 + i\mu R)e^{i\mu R} + (1 + \alpha R + \alpha^2 R^2)e^{-\alpha R}]. \quad (\text{A10b})$$

The Stokeslet also generates a rotational motion with an angular velocity of $(1/2)\nabla \times \hat{\mathbf{v}}_\omega$; therefore, the

translation-rotation cross-mobility is given by

$$\hat{\mu}_{12}^{rt}(\mathbf{R}, \omega) = -\frac{1}{2}\nabla \times \hat{\mathbf{G}}(\mathbf{R}, \omega), \quad (\text{A11})$$

which has a perpendicular component only:

$$\hat{\mu}_{12}^{rt\perp}(R, \omega) = \mu_0^r \frac{a^3}{R^2} (1 + \alpha R)e^{-\alpha R}. \quad (\text{A12})$$

There are no effects from the fluid compressibility. This result is also derived by considering the velocity field generated by a rotlet (point torque) [22]. The rotation-rotation cross-mobility is obtained from the angular velocity generated by a rotlet:

$$\hat{\mu}_{12}^{rr}(\mathbf{R}, \omega) = \frac{1}{4}\nabla \times \nabla \times \hat{\mathbf{G}}(\mathbf{R}, \omega). \quad (\text{A13})$$

Then, each component is given by

$$\hat{\mu}_{12}^{rr\parallel}(R, \omega) = \mu_0^r \frac{a^3}{R^3} (1 + \alpha R)e^{-\alpha R}, \quad (\text{A14a})$$

$$\hat{\mu}_{12}^{rr\perp}(R, \omega) = -\mu_0^r \frac{a^3}{2R^3} (1 + \alpha R + \alpha^2 R^2)e^{-\alpha R}. \quad (\text{A14b})$$

Using the analytical solutions for an isolated single particle for the self-mobilities and the Oseen approximation for the cross-mobilities, the velocity cross-relaxation functions can be approximated by Eq. (12).

-
- [1] W. van Saarloos and P. Mazur, *Physica A* **120**, 77 (1983).
 - [2] H. J. H. Clercx and P. P. J. M. Schram, *Physica A* **174**, 325 (1991).
 - [3] S. Henderson, S. Mitchell, and P. Bartlett, *Phys. Rev. Lett.* **88**, 088302 (2002).
 - [4] M. Atakhorrami, G. H. Koenderink, C. F. Schmidt, and F. C. MacKintosh, *Phys. Rev. Lett.* **95**, 208302 (2005).
 - [5] S. Martin, M. Reichert, H. Stark, and T. Gisler, *Phys. Rev. Lett.* **97**, 248301 (2006).
 - [6] Y. von Hansen, A. Mehlich, B. Pelz, M. Rief, and R. R. Netz, *Rev. Sci. Instrum.* **83**, 095116 (2012).
 - [7] P. Español, M. A. Rubio, and I. Zúñiga, *Phys. Rev. E* **51**, 803 (1995).
 - [8] P. Español, *Physica A* **214**, 185 (1995).
 - [9] Y. Nakayama and R. Yamamoto, *Phys. Rev. E* **71**, 036707 (2005).
 - [10] Y. Nakayama, K. Kim, and R. Yamamoto, *Eur. Phys. J. E* **26**, 361 (2008).
 - [11] R. Tatsumi and R. Yamamoto, *Phys. Rev. E* **85**, 066704 (2012).
 - [12] J. Happel and H. Brenner, *Low Reynolds Number Hydrodynamics*, (Kluwer Academic, Boston, 1983).
 - [13] D. J. Jeffrey and Y. Onishi, *J. Fluid Mech.* **139**, 261 (1984).
 - [14] M. Reichert and H. Stark, *Phys. Rev. E* **69**, 031407 (2004).
 - [15] J. D. Jackson, *Classical Electrodynamics* 3rd ed., (Wiley, New York, 1999).
 - [16] B. U. Felderhof, *Phys. Fluids* **19**, 073102 (2007).
 - [17] B. Cichocki and B. U. Felderhof, *Phys. Rev. E* **62**, 5383 (2000).
 - [18] D. Bedeaux and P. Mazur, *Physica* **78**, 505 (1974).
 - [19] B. U. Felderhof, *Physica A* **389**, 5602 (2010).
 - [20] B. U. Felderhof, *Phys. Fluids* **19**, 126101 (2007).
 - [21] B. U. Felderhof, *Physica A* **84**, 569 (1976).
 - [22] S. Kim and S. J. Karrila, *Microhydrodynamics*, (Butterworth-Heinemann, Boston, 1991).